



注意事項 Important Points

- 1. 當尿袋裝滿了2/3之尿液時, 必須把尿液倒掉。 Urine bag must be emptied when the urine bag filled more than 2/3 at the time.
- 2. 尿袋的位置必須常保持在低於膀胱之位置。以防逆行 件之尿道感染。
 - Keep the urine collection bag below the level of the bladder to prevent back flow of urine which will cause urinary tract infection.
- 3. 當尿袋破爛、或更換新導尿喉時,請更換尿袋。 Change the urine bag when the urine bag is broken, otherwise change it together with the foley catheter.
- 4. 檢查尿喉,避冤尿喉扭結,以防阻塞尿液流通。 Check the tubing is not kinked to avoid blocking of urine flow.
- 5. 一般導尿喉每一星期由醫護人員替病人更換。(矽膠 導尿喉每一個月更換一次)

Change the urine catheter by health care worker every week.

(Silione urine catheter change every month)

- 6. 鼓勵每天飲用6-8杯水,至少1.5公升,用肥皂和水 清潔尿道口,保持清潔衛生, 有助減少尿道感染。 To prevent urinary tract infection, use soap and water clean the urethral opening and drink 6-8 glasses of water daily (at least 1.5L/day).
- 7. 若有下列情況,請立即就醫求診: Call your health care provider immediately if you notice:
 - 發燒或發冷。 Fever or chills.
 - 膀胱赤痛。 Lower abdominal pain.
 - 小便帶血、混濁或有惡臭味。 Blood in urine, cloudy urine or strong odor.
 - 小便量少而膀胱仍感脹滿有尿意。 Minimal urinary output accompaning lower abdominal distention.
 - 當導尿管阻塞、脫出或有滲尿。 When the urine catheter have leakage, blockage or dislodged.







22 亞皆老街,九龍 Argyle Street, Kowloon **吞 2711 5222** www.evangel.org.hk





Caring for Your Urinary Catheter at Home



全人醫治 We provide holistic 盡心關懷 & compassionate care

家居護理尿袋 注意事項

Caring for Your Urinary Catheter at Home

所需物品 Materials:
70%消毒火酒棉塊及量杯(需要時)
70% alcohol pad and measuring jar
(when necessary)



