放回樽蓋並蓋好。

Replace and tighten the cap right away.



**刘**同時使用兩種不同的眼藥水或眼藥膏,應相隔 約十五分鐘。

If you are using two different eye drops or eye ointments, separate the doses by about 15 minutes.



## 注意 Note:

- 1. 眼藥水開蓋一個月後不可再使用。 Discard eye drops one month after first opening.
- 2. 切勿使用已變色或過期的眼藥水。
  Never use discoloured or expired eye drops.
- 3. 為防止傳播細菌及避免耽誤病情,切勿與他人共用 眼藥水。

Do not share your eye drops with others to avoid transmission of bacteria and delay in treatment.

4. 療程期間切勿佩戴隱形眼鏡。 Do not wear contact lens during treatment.

此藥物指引只供參考之用。

This information leaflet is for reference only.

用藥應按醫生指示。

Medicines should be used as directed by the doctor.

如有疑問請向主診醫生或藥劑師查詢。 For enquiry, please consult your doctor

For enquiry, please consult your doctor or pharmacist.

切勿將自己的藥物給予他人服用。
Do not share your medicine with others.









## 滴眼藥水使用方法

How to Use Eye Drops



## **滴眼藥水**使用方法 How to Use **Eye Drops**

用清水及肥皂徹底清潔雙手。
Wash your hands
thoroughly with
soap and
water.

2 將頭仰後,眼向上望,輕輕拉低下眼蓋成一袋形。
Tilt your head backwards, with your eyes looking upwards, gently pull down the lower eyelid to create a pouch.



 滴一滴眼藥水在眼袋内,切勿讓藥水樽滴咀接 觸到眼睛,眼蓋或雙手。 Instill one drop into the pouch. Avoid touching the tip of the bottle against your eye, eyelid or hands.



引上眼睛數分鐘,同時用手指輕壓眼頭。 Close your eyes for a few minutes. Apply gentle pressure to the inner corner of your eye.



用紙巾擦去流出眼外的藥水。
Wipe off any eye drops running out of your eye with a tissue.



如須要使用超過一滴眼藥水,應相隔五分鐘。
If you are to use more than one drop, wait for 5 minutes before instilling the next drop.

